

Self-consistent Approach to Off-Shell Transport*

Yu.B. Ivanov,^{1,†} J. Knoll,^{1,‡} and D.N. Voskresensky^{1,§}

¹*Gesellschaft für Schwerionenforschung mbH,*

Planckstr. 1, 64291 Darmstadt, Germany

The properties of two forms of the gradient expanded Kadanoff–Baym equations, i.e. the Kadanoff–Baym and Botermans–Malfliet forms, suitable to describe the transport dynamics of particles and resonances with broad spectral widths, are discussed in context of conservation laws, the definition of a kinetic entropy and the possibility of numerical realization. Recent results on exact conservations of charge and energy–momentum within Kadanoff–Baym form of quantum kinetics based on local coupling schemes are extended to two cases relevant in many applications. These concern the interaction via a finite range potential, and, relevant in nuclear and hadron physics, e.g. for the pion–nucleon interaction, the case of derivative coupling.

I. INTRODUCTION

Ever since L. Boltzmann had suggested his famous kinetic equation, the field of non-equilibrium physics and stochastic processes has grown tremendously, expanding into various directions. The interactions among particles driven by mean fields were included, quasiparticles were introduced in order to include much of the medium effects, the kinematics was extended to the relativistic case, ultimately theoretical foundations of transport equation were given from an underlying quantum many-body or field theory. In this line of achievements also stands the work of G.I. Budker and S.T. Belyaev who demonstrated the Lorentz invariance of the relativistic distribution function and derived relativistic Fokker–Planck kinetic equation [1]. The work entered into many textbooks and found numerous applications

* Dedicated to S.T. Belyaev on the occasion of his 80th birthday.

†Y.Ivanov@gsi.de; RRC “Kurchatov Institute”, Kurchatov sq. 1, Moscow 123182, Russia

‡J.Knoll@gsi.de

§D.Voskresensky@gsi.de; Moscow Institute for Physics and Engineering, Kashirskoe sh. 31, Moscow 115409, Russia

in atomic physics and electron-positron plasma. Presently the relativistic transport concepts are a conventional tool to analyze the dynamics of dense and highly excited matter produced in relativistic heavy-ion collisions.

Along with semiphenomenological extensions a great progress was achieved in microscopic foundation of the kinetic theory, which is mainly associated with the names of Bogolyubov, Born, Green, Kirkwood, Yvon, and Zubarev. The appropriate frame for description of non-equilibrium processes within the real-time formalism of quantum-field theory was developed by Schwinger, Kadanoff, Baym and Keldysh [2, 3, 4]. This formalism allows extensions of the kinetic picture beyond conventional approximations (like the quasiparticle one) and has found now numerous applications in many domains of physics.

The interest in transport descriptions of heavy-ion collisions beyond the quasiparticle approximation was initiated by Danielewicz [5], using the gradient expanded Kadanoff–Baym equations. These attempts have recently has been revived [6, 7, 8, 9, 10, 11, 12, 13] in order to properly describe the transport properties of broad resonances (like the ρ meson and Δ isobar). In the dense environment also stable particle acquire a considerable width because of collisional broadening. A proper dynamical treatment of their widths in the dense nuclear medium within a transport theoretical concepts is still a challenging problem. Transport approaches for treating such off-shell dynamics were proposed in refs. [7, 8, 9, 10, 11, 12, 13]. They all were based on the Kadanoff–Baym equations [3, 14] which describe the non-equilibrium quantum evolution at the truncation level of the Schwinger–Dyson equation. Expanded up to the first space–time gradients these provide equations for the one-body phase-space distribution functions with collision term and Poisson-bracket terms arising from the first order gradient terms. Presently two slightly different forms of the gradient-expanded Kadanoff–Baym equations are used: the original Kadanoff–Baym (KB) form [3], as it follows right after the gradient expansion without any further approximations, and the Botermans–Malfliet (BM) one [15], which is derived from the KB form by omitting certain second-order space-time gradient corrections.

In this paper we would like to compare these two forms of “quantum” kinetic equations and discuss their advantages and disadvantages from the point of view of their conserving properties, the possibility of numerical realization, etc., sect. II. Technical details on the conserving-properties are deferred to the appendices (B–D), since they illustrate some of the general consideration of ref. [10] together with some extensions to cases particularly relevant

in nuclear physics. Append. C treats nonrelativistic nucleon–nucleon interactions via a potential of finite range. The derivative coupling is considered in Append. D at the example of the P -wave pion–nucleon interaction. In sect. III, we supplement some considerations about the construction of a kinetic entropy within these two transport schemes. To make the paper self-contained, we summarize the time-contour matrix notation in Append. A, and introduce the Φ -functional formalism for derivative coupling in Append. B. A summary is given in sect. IV.

II. OFF-SHELL KB AND BM KINETICS

In this section we summarize the formulation of the off-shell kinetic equations in the two different forms: in the KB form, i.e. as it follows right after the gradient expansion of the exact Kadanoff–Baym equations, and in the BM form [15] which differs from the KB form only in the second order of the gradient expansion. We assume the reader to be familiar with the real-time formulation of non-equilibrium many-body theory and use the contour matrix notation, detailed in Append. A.

Starting point of all considerations is the set of Kadanoff–Baym equations which express the space-time changes of the Wigner transformed correlation function $iG^{-+}(X, p)$ in terms of the real-time contour convolution of the self-energy Σ with the Green function G . We give the kinetic equation in compact notation (cf. Eq. (A4))

$$v_\mu \partial_X^\mu iG^{-+}(X, p) = [\Sigma \otimes G - G \otimes \Sigma]_{X,p}^{-+} \quad \text{with} \quad v^\mu = \frac{\partial}{\partial p_\mu} G_0^{-1}(p), \quad (1)$$

where $G_0^{-1}(p)$ is the Fourier transform of the inverse free Green function

$$G_0^{-1}(p) = \begin{cases} p^2 - m^2 & \text{for relativistic bosons} \\ p_0 - \mathbf{p}^2/(2m) & \text{for non-rel. fermions or bosons.} \end{cases} \quad (2)$$

For a complete definition, Eq. (1) has to be supplemented with further equations, e.g., for the retarded Green function together with the retarded relations (A6). In addition to these equations the exact set of Kadanoff–Baym equations also includes the prototype of the mass-shell equation, which we also discuss below. If a system under consideration is only slightly spatially inhomogeneous and slowly evolving in time, a good approximation is provided by an expansion up to first order in space–time gradients. Then the main problem to arrive at

a proper kinetic equation consists in accurately disentangling the rather complicated r.h.s. of Eq. (1). This problem in the context of conserving approximations will be addressed here.

A. Φ -derivable approximations

In actual calculations one has to use certain approximations or truncation schemes to the exact non-equilibrium theory, which make conserving properties (such as charge and energy-momentum conservations) and thermodynamic consistency of the transport theory not evident. It was shown [8, 16, 17] that there exist a class of self-consistent approximations, called Φ derivable approximations, which are conserving at the expectation value level and at the same time thermodynamically consistent, i.e. they provide true Noether currents and a conserved energy-momentum tensor. In these schemes the self-consistent self-energies are generated from a functional $\Phi[G]$ through the following variational procedure [8]

$$-i\Sigma_{ik}(X, p) = \mp \frac{\delta i\Phi[G]}{\delta iG^{ki}(X, p)} \times \begin{cases} 2 & \text{for real fields} \\ 1 & \text{for complex fields} \end{cases}, \quad i, k \in \{-+\}. \quad (3)$$

The functional $\Phi[G]$ specifies the truncation scheme. It consists of a set of properly chosen closed two-particle irreducible diagrams, where lines denote the self-consistent propagator G , while vertices are bare. The functional variation with respect to G diagrammatically implies an opening of a propagator line of Φ .

Particular examples of Φ derivable approximations can be found in Appendices C and D, which consider applications of the general formalism to cases important in nuclear physics. The treatment of the pion-nucleon derivative coupling in Append. D requires the corresponding extension of the Φ derivable formalism, which has not been done up to now. Therefore, in Append. B we perform such an extension and derive the relevant modifications of the variational rules and the ensuing additional terms in the current and energy-momentum tensor expressions.

The conserving properties of these approximations are *exact* at the level of Kadanoff-Baym equations (1), while after the expansion up to the first space-time gradients they are generally expected to be only *approximately* fulfilled.

B. Physical notation

It is helpful to eliminate the imaginary factors inherent in the standard Green function formulation and introduce quantities which are real and, in the quasi-homogeneous limit, positive, with clear physical meaning, thereby. Thus instead of Green functions $G^{ij}(X, p)$ and self-energies $\Sigma^{ij}(X, p)$ with $i, j \in \{-+\}$ (see Append. A) in the Wigner representation we use the kinetic notation of ref. [8]. We define¹

$$\begin{aligned} F(X, p) &= A(X, p)f(X, p) = (\mp)\mathrm{i}G^{-+}(X, p), \\ \tilde{F}(X, p) &= A(X, p)[1 \mp f(X, p)] = \mathrm{i}G^{+-}(X, p), \end{aligned} \quad (4)$$

for the generalized Wigner functions F and $\tilde{F}$ and the corresponding *four*-phase-space distribution functions $f(X, p)$ and Fermi/Bose factors $[1 \mp f(X, p)]$. Here

$$A(X, p) \equiv -2\mathrm{Im} G^R(X, p) = \tilde{F} \pm F \quad (5)$$

is the spectral function, and G^R is the retarded propagator. The spectral function satisfies the sum rule

$$\int_0^\infty \frac{dp_0}{2\pi} A(X, p) = 1 \quad \text{for nonrelativ. particles,} \quad (6)$$

$$\int_{-\infty}^\infty \frac{dp_0}{2\pi} p_0 A(X, p) = 1 \quad \text{for relativ. bosons,} \quad (7)$$

which follows from the canonical equal-time (anti) commutation relations for (fermionic) bosonic field operators. Likewise the gain and loss rates of the collision integral are defined as

$$\Gamma^{\mathrm{in}}(X, p) = \mp \mathrm{i}\Sigma^{-+}(X, p), \quad \Gamma^{\mathrm{out}}(X, p) = \mathrm{i}\Sigma^{+-}(X, p) \quad (8)$$

with the damping width

$$\Gamma(X, p) \equiv -2\mathrm{Im} \Sigma^R(X, p) = \Gamma^{\mathrm{out}}(X, p) \pm \Gamma^{\mathrm{in}}(X, p), \quad (9)$$

where Σ^R is the retarded self-energy.

In terms of above kinetic notation, the gradient-expanded Kadanoff–Baym equations are reduced to equations for four real quantities: two equations for the real and imaginary parts

¹ Here and below the upper sign corresponds to fermions, while the lower sign, to bosons.

of the retarded Green function, while there are two equations for the phase-space occupation F : the KB kinetic equation and the prototype “mass-shell equation”. The latter doubling of equations reflects the well known redundancy of the Kadanoff–Baym equations. Before the gradient expansion both equations are completely identical. However, after the gradient expansion their interrelation is no longer obvious and deserves special care (see below).

The equations for the retarded propagator in first-order gradient approximation can be immediately solved with the result [3, 15]

$$G^R = \frac{1}{M(X, p) + i\Gamma(X, p)/2} \Rightarrow \begin{cases} A(X, p) = \frac{\Gamma(X, p)}{M^2(X, p) + \Gamma^2(X, p)/4} \\ \text{Re } G^R(X, p) = \frac{M(X, p)}{M^2(X, p) + \Gamma^2(X, p)/4} \end{cases} \quad (10)$$

with the “mass” function

$$M(X, p) = G_0^{-1}(p) - \text{Re } \Sigma^R(X, p). \quad (11)$$

Although the solution (10) is simply algebraic, it is valid up to first-order gradients.

C. Kadanoff–Baym form

In terms of above notation, the KB kinetic equation for F in the first-order gradient approximation takes the form

$$\mathcal{D}F(X, p) - \{\Gamma^{\text{in}}, \text{Re } G^R\} = C(X, p). \quad (12)$$

We denote this as the *quantum transport equation in the KB-choice*². Here the differential drift operator is defined as

$$\mathcal{D} = \left(v_\mu - \frac{\partial \text{Re } \Sigma^R}{\partial p^\mu} \right) \partial_X^\mu + \frac{\partial \text{Re } \Sigma^R}{\partial X^\mu} \frac{\partial}{\partial p_\mu}, \quad (13)$$

and $\{\dots, \dots\}$ denotes the four-dimensional Poisson bracket

$$\{f(X, p), \varphi(X, p)\} = \frac{\partial f}{\partial p^\mu} \frac{\partial \varphi}{\partial X_\mu} - \frac{\partial f}{\partial X^\mu} \frac{\partial \varphi}{\partial p_\mu}, \quad (14)$$

² If the system consists of several different particle species, there is a set of coupled kinetic equations corresponding to each species (e.g., see Append. D).

in covariant notation. Please note that now after gradient approximation all quantities on the l.h.s. are to be taken in local approximation, i.e. void of any further gradient terms. Thus, the occurring self-energies are obtained evaluating the diagrams as in momentum representation with the coordinates X of all Greens functions kept identical. The r.h.s. specifies the collision term³

$$C(X, p) = \Gamma^{\text{in}}(X, p) \tilde{F}(X, p) - \Gamma^{\text{out}}(X, p) F(X, p). \quad (15)$$

If the diagrams for the self-energy contain internal vertices, which give rise to memory or non-local effects, the gain and loss rates contain additional gradient terms which have to be constructed, e.g., according to the rules given in [10]. The resulting local part of the collision term is charge (e.g., the baryonic number) and energy-momentum conserving by itself

$$\text{Tr} \int \frac{d^4 p}{(2\pi)^4} \begin{pmatrix} e \\ p^\mu \end{pmatrix} C^{\text{loc}} = 0. \quad (16)$$

Here and below, e denotes the elementary charge, while Tr implies the sum over all possible internal degrees of freedom, like spin, and over possible particle species. We do not explicitly introduce the particle-species label to avoid overcomplication of equations. In terms of a local functional Φ^{loc} the explicit form of the local collision term is

$$C^{\text{loc}}(X, p) = \frac{\delta i\Phi^{\text{loc}}}{\delta \tilde{F}(X, p)} \tilde{F}(X, p) - \frac{\delta i\Phi^{\text{loc}}}{\delta F(X, p)} F(X, p), \quad (17)$$

cf. Eq. (3). In this paper we limit the considerations to cases void of memory effects in this collision term. The latter effects were studied in [8].

Relation (16) permits us to derive the current

$$j_{\text{KB-eff}}^\mu(X) = e \text{Tr} \int \frac{d^4 p}{(2\pi)^4} \left[v^\mu F(X, p) + \text{Re} \Sigma^R \frac{\partial F}{\partial p_\mu} - \text{Re} G^R \frac{\partial \Gamma^{\text{in}}}{\partial p_\mu} \right]. \quad (18)$$

of a charge e (e.g., the baryonic one) from the KB kinetic equation (12) which is conserved

$$\partial_\mu j_{\text{KB-eff}}^\mu(X) = 0. \quad (19)$$

Note that this current formally differs from the true Noether current

$$j^\mu(X) = e \text{Tr} \int \frac{d^4 p}{(2\pi)^4} v^\mu F(X, p) + j_{(\text{der})}^\mu(X), \quad (20)$$

³ See an example in Eq. (C3).

which follows right from the operator expression for this quantity, cf. ref. [8] and Appendix B. The additional term $j_{(\text{der})}^\mu$ appears only in the case of derivative coupling, see Eq. (B19). In view of the gradient approximation employed, one could generally expect both currents to differ beyond the validity range of the gradient approximation. However, as demonstrated in detail in ref. [10], these two currents are exactly equal for Φ -derivable approximations, if a consistent gradient expansion is performed also in the gain and loss rates (8) of the collision term (15). In this case the exact conservation of the Noether current results from the corresponding invariance of the Φ functional – (Eq. (6.9) in [17]) – which survives the gradient expansion

$$e\text{Tr} \int \frac{d^4p}{(2\pi)^4} [\{\text{Re} \Sigma^R, F\} - \{\text{Re} G^R, \Gamma^{\text{in}}\} + C] = \partial_\mu j_{(\text{der})}^\mu. \quad (21)$$

The latter relation is written down for the general case of memory or non-local effects included in C . If such effects in C are absent, the collision term drops out of Eq. (21) according to Eq. (16).

Within Φ -derivable approximations also the conservation of energy–momentum can be established, for local (point-like) couplings providing a local energy-momentum tensor. The p^ν weighted four-momentum integral of the KB kinetic equation leads to the following consistency relation [8]⁴

$$\begin{aligned} & \partial^\nu (\mathcal{E}^{\text{pot}} - \mathcal{E}^{\text{int}}) - \partial_\mu \mathcal{E}_{(\text{der.})}^{\mu\nu} \\ &= \text{Tr} \left(\frac{1}{2} \right)_{\text{n.b.}} \int \frac{p^\nu d^4p}{(2\pi)^4} [\{\text{Re} \Sigma^R, F\} - \{\text{Re} G^R, \Gamma^{\text{in}}\} + C], \end{aligned} \quad (22)$$

which is again *exact* after the gradient expansion as shown in ref. [10] (see also Appendices C and D). It implies that the Noether energy–momentum tensor

$$\Theta^{\mu\nu}(X) = \text{Tr} \left(\frac{1}{2} \right)_{\text{n.b.}} \int \frac{d^4p}{(2\pi)^4} v^\mu p^\nu F(X, p) + g^{\mu\nu} (\mathcal{E}^{\text{int}} - \mathcal{E}^{\text{pot}}) + \mathcal{E}_{(\text{der.})}^{\mu\nu} \quad (23)$$

is *exactly* conserved by the kinetic equation (12)

$$\partial_\mu \Theta^{\mu\nu}(X) = 0. \quad (24)$$

Here potential energy density $\mathcal{E}^{\text{pot}}(X)$, which a probe particle with Wigner density $F(X, p)$ would experience due to the interaction with all other particles in the system, is

$$\mathcal{E}^{\text{pot}}(X) = \text{Tr} \left(\frac{1}{2} \right)_{\text{n.b.}} \int \frac{d^4p}{(2\pi)^4} [\text{Re} \Sigma^R F + \text{Re} G^R \Gamma^{\text{in}}]. \quad (25)$$

⁴ Here in compliance with Eq. (3) we define the factor $\left(\frac{1}{2} \right)_{\text{n.b.}} = \begin{cases} 1/2 & \text{for neutral bosons (real fields)} \\ 1 & \text{else} \end{cases}$.

The interaction energy density $\mathcal{E}^{\text{int}}(X)$ specifies that part of the total energy density which is due to interactions. In simple cases it relates to $\mathcal{E}^{\text{pot}}$ by a simple counting factor, namely, if all the interaction vertices of a theory have the same number n_l of lines attached to them

$$\mathcal{E}^{\text{int}}(X) = \frac{2}{n_l} \mathcal{E}^{\text{pot}}(X). \quad (26)$$

In particular, for two-body interactions one has $n_l = 4$ and thus $\mathcal{E}^{\text{int}} = \frac{1}{2} \mathcal{E}^{\text{pot}}$, while for the fermion–boson interaction $n_l = 3$, which results in $\mathcal{E}^{\text{int}} = \frac{2}{3} \mathcal{E}^{\text{pot}}$. In Appendices C and D we discuss cases of this type. The additional term $\mathcal{E}_{(\text{der.})}^{\mu\nu}$ appears in Eq. (23) only in the case of derivative coupling, cf. Eq. (B18).

The considerations given above summarize the results of ref. [10] which are quite general. However they are restricted to local (point-like) interactions and void of derivative couplings. This excludes two important cases relevant to many areas in physics, nuclear physics in particular. These are the cases of interaction mediated by finite range non-relativistic potentials and of derivative couplings like the P -wave pion–nucleon interaction. Since the considerations are rather technical they are exemplified in Appendices C and D. There the results of [10] are generalized proving that also in these cases conserved currents and expressions for a conserved total energy and total momentum can be constructed. These two appendices, also provide further illustrations of the discussion given in the present section.

The conserving feature are especially important for devising numerical simulation codes based on this kinetic equation. Indeed, if a test-particle method is used, one should be sure that the number of test particles is exactly conserved rather than approximately. For a direct application of this method, however, there is a particular problem with the KB kinetic equation. In the test-particle method the distribution functions are represented by an ensemble of test particles as follows

$$F(X, p) \sim \sum_i \delta^{(3)}(\mathbf{X} - \mathbf{X}_i(T)) \delta^{(4)}(p - p_i(T)), \quad (27)$$

where the i -sum runs over test particles. Then the $\mathcal{D}F$ term in Eq. (12) just corresponds to the classical motion of these test particles subjected to forces inferred from $\text{Re } \Sigma^R$, while the collision term C gives stochastic change of test-particles' momenta, when their trajectories “cross”. The additional term, i.e. the Poisson-bracket term $\{\Gamma^{\text{in}}, \text{Re } G^R\}$, spoils this simplistic picture, since derivatives acting on the distribution function F appear here only indirectly. Namely they are encoded through derivatives of Γ^{in} . This term is responsible

for backflow effects which restore the Noether current to be the conserved one. However, such backflow phenomena are difficult to absorb into test particles, since they describe the response of the medium to the motion of the charges. In order to conserve the number of test particles between subsequent collisions, one would have to unite the additional term $\{\Gamma^{\text{in}}, \text{Re } G^R\}$ with the drift term $\mathcal{D}F$ even in the simplest case, when the collision term is charge conserving by itself (see Eq.(16)) and derivative currents vanish, $j_{(\text{der})}^\mu = 0$. However, the interpretation of the additional term $\{\Gamma^{\text{in}}, \text{Re } G^R\}$ causes problems within this picture, since it is not just proportional to the same δ -functions as in Eq. (27) and thus can not be included in the collisionless propagation of test particles. This problem, of course, does not prevent a direct solution of the kinetic equation. E.g., one can apply well developed lattice methods, which are, however, much more complicated and time-consuming as compared to the test-particles approach.

Within the same approximation level the set of Dyson equations for Green functions $G^{ij}(X, p)$ provides us with an alternative equation for F

$$MF - \text{Re } G^R \Gamma^{\text{in}} = \frac{1}{4} (\{\Gamma, F\} - \{\Gamma^{\text{in}}, A\}), \quad (28)$$

which is called the mass-shell equation, since in the quasiparticle limit it provides the mass-condition $M = 0$. This equation coincides with the kinetic one (12) only within the first-order gradient approximation [8, 11, 12, 15], while both equations are *exactly* identical before the gradient expansion. In view of this still remaining difference the practical recipe is to forget about the mass-shell equation (28), since the retarded equation (10) determines the spectral distribution, and to treat Eq. (12) as a proper quantum kinetic equation. Still this is an ambiguous recipe, which historically was one of the motivation to proceed to the Botermans–Malfliet form of the quantum kinetic equation.

D. Botermans–Malfliet form

As can be seen from the mass-shell equation (28) and Eq. (12) [8, 11, 12, 15], the gain rate Γ^{in} departs from $F\Gamma/A$ only by corrections of first order in the gradients

$$\Gamma^{\text{in}} = \Gamma F/A + O(\partial_X) \quad (29)$$

(in equilibrium both equate each another). This fact permits to substitute the r.h.s. estimate for Γ^{in} in any of the gradient terms, i.e. in the Poisson-bracketed terms of Eqs. (12) and

(28) and neglect the correction $O(\partial_X)$ as it leads to terms of already second-order in the gradients. Upon this substitution, first proposed by Botermans and Malfliet [15], one arrives at the following form of the kinetic and mass-shell equations

$$\mathcal{D}F(X, p) - \left\{ \Gamma \frac{F}{A}, \text{Re } G^R \right\} = C(X, p), \quad (30)$$

$$MF - \text{Re } G^R \Gamma^{\text{in}} = \frac{1}{4} \left(\{ \Gamma, F \} - \left\{ \frac{\Gamma F}{A}, A \right\} \right), \quad (31)$$

which are already *exactly* identical, as they were before the gradient expansion, and still equivalent to those in the KB form within the first-order gradient approximation. The so obtained equation (30) will be called *quantum kinetic equations in BM-choice*. This equation *exactly* conserves the following effective current

$$j_{\text{BM-eff}}^\mu(X) = e \text{Tr} \int \frac{d^4 p}{(2\pi)^4} \left[v^\mu F(X, p) + \text{Re } \Sigma^R \frac{\partial F}{\partial p_\mu} - \text{Re } G^R \frac{\partial(\Gamma F/A)}{\partial p_\mu} \right], \quad (32)$$

which differs from the Noether current j^μ in terms of the order of $O(\partial_X)$, provided a Φ -derivable approximation is used for self-energies. All KB-choice properties of Eq. (30) within a Φ -derivable approximation also transcribe to BM-choice through the substitution $\Gamma^{\text{in}} = \Gamma F/A$ in Eqs. (21), (22) and (25). This substitution, however, touches the accuracy of those relations. For instance, the conservation laws of the Noether currents (20) and the energy-momentum tensor (23) are then only approximately fulfilled together with the corresponding consistency relations (21) and (22) which now look as

$$e \text{Tr} \int \frac{d^4 p}{(2\pi)^4} [\{ \text{Re } \Sigma^R, F \} - \{ \text{Re } G^R, \Gamma F/A \} + C] \simeq \partial_\mu j_{(\text{der})}^\mu, \quad (33)$$

$$\begin{aligned} & \partial^\nu (\mathcal{E}^{\text{pot}} - \mathcal{E}^{\text{int}}) - \partial_\mu \mathcal{E}_{(\text{der.})}^{\mu\nu} \\ & \simeq \text{Tr} \left(\frac{1}{2} \right)_{\text{n.b.}} \int \frac{p^\nu d^4 p}{(2\pi)^4} \left[\{ \text{Re } \Sigma^R, F \} - \left\{ \text{Re } G^R, \frac{\Gamma F}{A} \right\} + C \right], \end{aligned} \quad (34)$$

respectively, and hold only up to first-order gradients.

The effective BM-current (32) was used by S. Leupold [12] as a basis for the construction of a test-particle ansatz for the nonrelativistic case. In this case the additional term $\{ \Gamma F/A, \text{Re } G^R \}$ in the BM kinetic equation (30) is expressed in terms of the same distribution function as the drift term $\mathcal{D}F$. Therefore, one can unify these terms to construct

equations of motions for test particles, which provide exact conservation of $j_{\text{BM-eff}}^\mu$. To automatically fulfill this effective current conservation, the test-particle ansatz is introduced for the combination

$$\frac{1}{2}\Gamma A \left(1 - \frac{\partial \text{Re } \Sigma^R}{\partial p_0} - \frac{M}{\Gamma} \frac{\partial \Gamma}{\partial p_0}\right) F(X, p) \sim \sum_i \delta^{(3)}(\mathbf{X} - \mathbf{X}_i(T)) \delta^{(4)}(p - p_i(T)), \quad (35)$$

rather than for the distribution function itself. Note that the energy $p_i^0(T)$ of the test particle is a free coordinate, not restricted by a mass-shell condition. W. Cassing and S. Juchem [11] extended this test-particle ansatz to the relativistic case. The equations of motion for the test particle, which follow from this ansatz, in particular give the time evolution of the off-shellness of a test particle [11, 12]

$$\frac{dM}{dT} = \frac{M}{\Gamma} \frac{d\Gamma}{dT}, \quad (36)$$

the origin of which can be traced back to the additional term $\{\Gamma F/A, \text{Re } G^R\}$ in the BM kinetic equation (30). Here M is the mass of the test particle relative to its on-shell value, see Eq. (11), and this equation of motion implies that once the width drops in time the particles are driven towards the on-shell mass $M = 0$. This clarifies the meaning of the latter term for the off-shell BM transport: it provides the time evolution of the off-shellness.

III. ENTROPY

Another important feature of the kinetic description is the approach to thermal equilibrium during evolution of a closed system. In terms of transport theory the sufficient (while not necessary!) condition of it is the existence of an H-theorem. Leaving aside all complications associated with non-local effects in the collision term and possible lack of positive definiteness of the transition rates, discussed in ref. [8], we confine our consideration to simple approximations, cf. (C1). As demonstrated in ref. [8], in the BM approximation to the quantum kinetic equations (30) the H-theorem can indeed be formulated

$$\partial_\mu s_{\text{BM}}^\mu(X) = \text{Tr} \int \frac{d^4 p}{(2\pi)^4} \ln \frac{\tilde{F}}{F} C^{\text{loc}}(X, p) \geq 0, \quad (37)$$

where the quantity

$$s_{\text{BM}}^\mu = \text{Tr} \int \frac{d^4 p}{(2\pi)^4} \left[\left(v^\mu - \frac{\partial \text{Re } \Sigma^R}{\partial p_\mu} \right) \left(\mp \tilde{F} \ln \frac{\tilde{F}}{A} - F \ln \frac{F}{A} \right) \right]$$

$$- \operatorname{Re} G^R \left(\mp \ln \frac{\tilde{F}}{A} \frac{\partial}{\partial p_\mu} \left(\Gamma \frac{\tilde{F}}{A} \right) - \ln \frac{F}{A} \frac{\partial}{\partial p_\mu} \left(\Gamma \frac{F}{A} \right) \right) \right] \quad (38)$$

obtained from the l.h.s. of the BM kinetic equation (30) is interpreted as a entropy flow for the BM-choice. For the Φ -derivable approximation (C1) the r.h.s. of relation (37) takes the following form

$$\begin{aligned} \operatorname{Tr} \int \frac{d^4 p}{(2\pi)^4} \ln \frac{\tilde{F}}{F} C^{\text{loc}}(X, p) &= \operatorname{Tr} \frac{1}{4} \int \frac{d^4 p_1}{(2\pi)^4} \frac{d^4 p_2}{(2\pi)^4} \frac{d^4 p_3}{(2\pi)^4} \frac{d^4 p_4}{(2\pi)^4} \\ &\times R^{\text{loc}} (2\pi)^4 \delta^4(p_1 + p_2 - p_3 - p_4) \left(F_1 F_2 \tilde{F}_3 \tilde{F}_4 - \tilde{F}_1 \tilde{F}_2 F_3 F_4 \right) \ln \frac{F_1 F_2 \tilde{F}_3 \tilde{F}_4}{\tilde{F}_1 \tilde{F}_2 F_3 F_4}, \end{aligned} \quad (39)$$

where R^{loc} is the transition rate determined by Eq. (C4). This expression is indeed non-negative, since $(x-y) \ln(x/y) \geq 0$ for any positive x and y , and is of the second order in deviation from equilibrium $(F - F_{\text{eq}})$, as both $(F_1 F_2 \tilde{F}_3 \tilde{F}_4 - \tilde{F}_1 \tilde{F}_2 F_3 F_4)$ and $\ln(F_1 F_2 \tilde{F}_3 \tilde{F}_4 / \tilde{F}_1 \tilde{F}_2 F_3 F_4)$ approach zero at equilibrium. From the kinetic equation it follows that the deviation from equilibrium is of the first order in time gradients: $(F - F_{\text{eq}}) \propto O(\partial_T F)$. This implies that the r.h.s. of relation (37) is of the second order in time gradients, which is, strictly speaking, beyond our first-order gradient approximation. However, from the point of view of practical use this feature is highly welcome as it guarantees equilibration. A further advantaged of the kinetic entropy flux (38) is that in equilibrium its entropy merges that thermodynamic expression deduced from the thermodynamic potential in the Φ derivable scheme [8, 18, 19].

In the case of the KB-choice (12) the situation is more controversial. Performing all the same manipulations with the KB kinetic equation (12) as those in ref. [8], we arrive at the following relation

$$\partial_\mu s_{\text{KB}}^\mu(X) = \operatorname{Tr} \int \frac{d^4 p}{(2\pi)^4} \ln \frac{\tilde{F}}{F} C^{\text{loc}} - \delta H_{\text{KB}}, \quad (40)$$

where

$$\begin{aligned} s_{\text{KB}}^\mu &= \operatorname{Tr} \int \frac{d^4 p}{(2\pi)^4} \left[\left(v^\mu - \frac{\partial \operatorname{Re} \Sigma^R}{\partial p_\mu} \right) \left(\mp \tilde{F} \ln \frac{\tilde{F}}{A} - F \ln \frac{F}{A} \right) \right. \\ &\quad \left. - \operatorname{Re} G^R \left(\mp \ln \frac{\tilde{F}}{A} \frac{\partial \Gamma^{\text{out}}}{\partial p_\mu} - \ln \frac{F}{A} \frac{\partial \Gamma^{\text{in}}}{\partial p_\mu} \right) \right]. \end{aligned} \quad (41)$$

$$\delta H_{\text{KB}} = - \int \frac{d^4 p}{(2\pi)^4} \operatorname{Re} G^R \left\{ \ln \frac{\tilde{F}}{F}, \frac{C^{\text{loc}}}{A} \right\}. \quad (42)$$

The KB entropy flow s_{KB}^μ is identical to the BM one s_{BM}^μ up to zero-order gradients, while they differ in the first-order gradient corrections. One can easily obtain s_{BM}^μ from the KB entropy flow by doing replacement (29) in Γ^{in} and a similar replacement in Γ^{out} .

The additional term δH_{KB} on the r.h.s. of relation (40) is of the second order in gradients, due to the Poisson bracket and $C^{\text{loc}} \propto O(\partial_T F)$. Therefore, the r.h.s. of (40) consists of two terms, which are of the same order of magnitude and one of them (δH_{KB}) is sign indefinite. This prevents us from concluding the positive definiteness of the r.h.s. of Eq. (40). Alternatively we were not able to cast this term into a full divergence as to be included into the definition of the KB entropy flow. This fact by itself does not imply that the system does not approach equilibrium or even the absence of an H-theorem for the KB kinetic equation but suggests that equilibration should be tested in actual calculations. The local H-theorem we are looking for is a very stringent condition, providing monotonous approach to equilibrium. In fact, equilibration may well be nonmonotonous in time.

Still, for the KB kinetic equation we are able to prove the H-theorem in a limiting case, i.e. close to local thermal equilibrium or for quasi-stationary state, which slowly evolve in space and time. To be definite, let us talk about the local thermal equilibrium. In terms of the distribution function

$$F(X, p) = F_{\text{loc.eq.}}(X, p) + \delta F(X, p) \quad (43)$$

the above assumption implies that $|\delta F| \ll F_{\text{loc.eq.}}$ and $|\partial_X F_{\text{loc.eq.}}| \lesssim |\partial_X \delta F|$. Then we can write down

$$\delta H_{\text{KB}} = \partial_\mu \delta s_{\text{KB}}^\mu(X) + \int \frac{d^4 p}{(2\pi)^4} \frac{C^{\text{loc}}}{A} \left\{ \ln \frac{\tilde{F}}{F}, \text{Re } G^R \right\}. \quad (44)$$

where

$$\delta s_{\text{KB}}^\mu(X) = -\text{Tr} \int \frac{d^4 p}{(2\pi)^4} \frac{\text{Re } G^R}{A} \frac{\partial \ln(\tilde{F}/F)}{\partial p_\mu} C^{\text{loc}}(X, p). \quad (45)$$

Here, the remaining term

$$\int \frac{d^4 p}{(2\pi)^4} \frac{C^{\text{loc}}}{A} \left\{ \ln \frac{\tilde{F}}{F}, \text{Re } G^R \right\} \propto O(\delta F \partial_X \delta F) + O(\delta F \partial_X F_{\text{loc.eq.}}), \quad (46)$$

can be neglected, as it has additional gradient smallness as compared to the first term on the r.h.s. of Eq. (40). Here, we have taken into account that $C^{\text{loc}} \propto \delta F$ and

$\left\{\ln(\tilde{F}/F), \text{Re } G^R\right\} \propto \partial_X(F_{\text{loc.eq.}} + \delta F)$. Thus, from Eq. (40) we conclude that

$$\partial_\mu (s_{\text{KB}}^\mu + \delta s_{\text{KB}}^\mu) \geq 0 \quad \text{near local equilibrium,} \quad (47)$$

which is the H-theorem for this case with the total entropy flow $s_{\text{KB}}^\mu + \delta s_{\text{KB}}^\mu$. Note that δs_{KB}^μ is proportional to the collision term and hence equals zero in equilibrium. The applicability range of this result is the same as that for the memory entropy derived in [8] for the BM-choice.

IV. SUMMARY AND PERSPECTIVES

In conclusion, we would like to summarize the present status of the two considered approaches to off-shell transport.

From a consistency point of view, the BM-choice looks more appealing, since it preserves the *exact* identity between the kinetic and mass-shell equation, a property inherent in the original KB equations [8]. For the KB-choice this identity between the kinetic and mass-shell equation is only *approximately* preserved, namely within the validity range of the first-order gradient approximation. However, this disadvantage is not of great practical use, since, in any case, only one of these two equations, namely the kinetic one, should be used in actual calculations.

For the construction of conservation laws related to global symmetries or energy and momentum the *local* collision term entirely drops out of the balance. Thus, the conservation laws solely depend on the properties of the first order gradient terms in the kinetic equation. In this respect the KB kinetic equation has a conceptual advantage as it leads to *exact* [10] rather than approximate conservation laws, provided the scheme is based on Φ -derivable approximations. Thereby the expectation values of the original operator expressions of conserved quantities (e.g., Noether currents) are exactly conserved. The reason is that the KB kinetic equation preserves certain contour symmetries among the various gradient terms, while they are violated for the BM-choice. Of course, within their range of applicability these two approaches are equivalent, because the BM kinetic equation conserves the charge and energy-momentum within the theoretical accuracy of the gradient approximation. Still, the fact that the KB-choice possesses exact conservation laws put this version to the level of a generic equation, much like the Boltzmann or hydrodynamic equations, to be used as phe-

nomenological dynamical equations for practical applications. Such conserving dynamical schemes may be useful even though the applicability condition of the approximation might be violated at some stages of evolution. For instance, such a situation happens at the initial stage of heavy-ion collisions. As the conservations are exact, we can still use the gradient approximation, relying on a minor role of this rather short initial stage in the total evolution of a system. Moreover, exact conservation laws allow us to keep control of numerical codes.

Although the KB kinetic equations posses exactly conserved Noether currents, a practical numerical approach (e.g., by a test-particle method) for its solution has not yet been established. The obstacle is the special Poisson-bracket term in the KB kinetic Eq. (12) which lacks proper interpretation since the phase-space occupation function $F(X, p)$ enters only indirectly through the gain-rate gradient terms. What is known is that this term encodes the backflow component which ensures the Noether currents to be conserved. However such backflow features are difficult to be implemented into a test-particle scheme. This problem, of course, does not exclude solution of the KB kinetic equation, e.g within well adapted lattice methods, which are, however, much more complicated and time-consuming as compared to the test-particle approach. For the BM kinetic equation, on the other hand, an efficient test-particle method is already available [11, 12], for the price that it deals with an alternative current rather than the Noether current.

As a novel part we showed (cf. appendices) that the exact conservation laws in the KB kinetic equations, originally derived for local interaction terms which lead to a local energy momentum tensor, also do hold for derivative couplings and for interactions of finite range, like a non-relativistic potential. For the latter case, of course, only global conservation of energy and momentum can be achieved. In order to deal with the derivative coupling, we extended the Φ -derivable approach to this case.

An important feature of kinetic descriptions is the approach to thermal equilibrium during evolution of a closed system. A sufficient (while not necessary!) condition is provided by an H-theorem. As was demonstrated in ref. [8], at least, within simplest Φ -derivable approximations for the kinetic equation in BM-choice an H-theorem indeed can be proven. The so derived kinetic entropy merged the equilibrium expression which in the context of Φ -derivable approximations results from the thermodynamic potential, cf. [8, 18, 19]. For the KB kinetic equation the result is by far weaker. Here we were able to prove the H-theorem only within simplest Φ -derivable approximations and for a system very close to almost spa-

tially homogeneous thermal local equilibrium or stationary state. These results, in general, do not imply that the system does not approach equilibrium but suggests that equilibration should be tested in actual calculations. Furthermore, the local H-theorem with a local entropy current which we considered for the BM-case may be by far too restrictive, providing monotonous approaching to equilibrium. In fact, for kinetic equations with memory or non-local effects equilibration may well be nonmonotonous in time.

Though the discussion in this paper is confined to problems of Φ -derivable off-shell transport based on the first-order gradient expansion, essential progress has recently been achieved also in solving KB equations directly without any gradient expansion for selected examples. These concern non-equilibrium processes in scalar and spinor–scalar models on 1 and 3 space dimensions, see e.g. [20, 21, 22] and references therein. It was found [20, 21] that after a comparably short but violent non-equilibrium evolution the time dependence of the Wigner transformed spectral function becomes rather weak even for moderate coupling constants. During this slow evolution the system is still far away from equilibrium. This fact provides a necessary condition for a successful gradient expansion and hence indicates a wide range of applicability of the approaches discussed in this paper. Even though the rapid far-from-equilibrium dynamics is formally beyond the scope of applicability of the gradient-expanded quantum kinetics, nevertheless, the KB choice includes all the ingredients required for such a treatment, i.e. the proper mean-field dynamics, together with the off-shell transport of particles, thereby satisfying exact rather than approximate conservation laws even far away from equilibrium.

Further progress in understanding the properties of Φ -derivable approximations to finite-temperature quantum field theory was reported concerning the question of renormalizability. The new results are equally applicable to quantum kinetic equations, both in KB or BM choices. In ref. [23] it was shown that truncated non-perturbative self-consistent Dyson resummation schemes can be renormalized with local counter terms defined at the vacuum level. The requirements are that the underlying theory is renormalizable and that the self-consistent scheme follows Baym’s Φ -derivable concept. This result proves that there is no arbitrariness in studying the in-medium modifications of model parameters like the mass and the coupling constants within this class of approximation schemes. It is sufficient to adjust them in the vacuum, for instance, by fitting them to scattering data, in order to predict their changes in medium without ambiguity. This result also guarantees the standard Φ -

derivable properties like thermodynamic consistency and exact conservation laws also for the renormalized approximation schemes to hold. In ref. [24] the theoretical concepts for the renormalization, devised in [23], were applied to the ϕ^4 -model, demonstrating the practicability of the method.

In general, the symmetries of the classical action which lead to Ward-Takahashi identities for the proper vertex-functions are violated for Φ -derivable approximations for functions beyond the one-point level, i.e., on the correlator level. This causes problems concerning the Nambu–Goldstone modes [25] in the broken symmetry case or concerning local symmetries (gauge symmetries) [26] on a level where the gauge fields are treated beyond the classical field approximation, i.e. on the propagator level. In ref. [26] it was shown that on top of any solution of a Φ -derivable approximation which is constructed from a symmetric Lagrangian there exists a non-perturbative effective action which generates proper vertex functions in the same sense as the 1PI effective action. These external vertex functions fulfill the Ward–Takahashi identities of the underlying symmetry. However, in general they coincide with the self-consistent ones only up to one-point order. Thus, usually the so generated external self-energy and higher vertex functions are different from the Φ -derivable expressions. Therefore, the pleasant property of the Φ -derivable approximations, namely the conserving one, proves to be lost. The derivation of approximation schemes that fulfill all symmetry properties of the underlying classical action and at the same time are fully self-consistent and conserving still remains as an open task.

As has been already mentioned, the gauge invariance may be lost in Φ -derivable approximations, too. In particular, this problem prevents applications of Φ -derivable approximations (including kinetic ones) to description quark–gluon plasma, based on QCD. This occurs because, in general, solution for dressed propagators and vertices do not satisfy Ward–Takahashi identities. This pathology shows up as an explicit dependence of results on the choice of the gauge condition. In ref. [27] it was demonstrated, in fact, that Φ -derivable approximations have a controlled gauge dependence, i.e. the gauge dependent terms appear at orders higher than the truncation order. Furthermore, using the stationary point obtained for the approximation to evaluate the complete 2PI effective action boosts the order, at which the gauge dependent terms appear, to twice the order of truncation. This is still not a solution of the gauge problem in the rigorous sense but certain progress to its better control and understanding.

Acknowledgments

We are grateful to G. Baym, J. Berges, P. Danielewicz, H. Feldmeier, B. Friman, H. van Hees, C. Greiner, E.E. Kolomeitsev and S. Leupold for fruitful discussions on various aspects of this research. This work was supported in part by the Deutsche Forschungsgemeinschaft (DFG project 436 RUS 113/558/0-2), the Russian Foundation for Basic Research (RFBR grants 03-02-04008 and 00-15-96590) and the German BMBF (contract RUS-01/690).

Appendix A: MATRIX NOTATION

In calculations that apply the Wigner transformations, it is necessary to decompose the full contour into its two branches—the *time-ordered* and *anti-time-ordered* branches. One then has to distinguish between the physical space-time coordinates $x, \dots$ and the corresponding contour coordinates x^c which for a given x take two values $x^- = (x_\mu^-)$ and $x^+ = (x_\mu^+)$ ($\mu \in \{0, 1, 2, 3\}$) on the two branches of the contour (see figure 1).




Figure 1: Closed real-time contour with two external points x, y on the contour.

Closed real-time contour integrations can then be decomposed as

$$\int_{\mathcal{C}} dx \dots = \int_{t_0}^{\infty} dx \dots + \int_{\infty}^{t_0} dx \dots = \int_{t_0}^{\infty} dx \dots - \int_{t_0}^{\infty} dx \dots, \quad (\text{A1})$$

where only the time limits are explicitly given. The extra minus sign of the anti-time-ordered branch can conveniently be formulated by a $\{-+\}$ “metric” with the metric tensor in $\{-+\}$ indices

$$(\sigma^{ij}) = (\sigma_{ij}) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (\text{A2})$$

which provides a proper matrix algebra for multi-point functions on the contour with “co”- and “contra”-contour values. Thus, for any two-point function F , the contour values are defined as

$$F^{ij}(x, y) := F(x^i, y^j), \quad i, j \in \{-, +\}, \quad \text{with}$$

$$\begin{aligned}
F_i^j(x, y) &:= \sigma_{ik} F^{kj}(x, y), & F_j^i(x, y) &:= F^{ik}(x, y) \sigma_{kj} \\
F_{ij}(x, y) &:= \sigma_{ik} \sigma_{jl} F^{kl}(x, y), & \sigma_i^k &= \delta_{ik}
\end{aligned} \tag{A3}$$

on the different branches of the contour. Here summation over repeated indices is implied. Then contour folding of contour two-point functions, e.g., in Dyson equations, simply becomes

$$\begin{aligned}
H(x^i, y^k) &= H^{ik}(x, y) = [F \otimes G]^{ik} \\
&\equiv \int_{\mathcal{C}} dz F(x^i, z) G(z, y^k) = \int dz F_j^i(x, z) G^{jk}(z, y)
\end{aligned} \tag{A4}$$

in the matrix notation.

For any multi-point function the external point x_{max} , which has the largest physical time, can be placed on either branch of the contour without changing the value, since the contour-time evolution from x_{max}^- to x_{max}^+ provides unity. Therefore, one-point functions have the same value on both sides on the contour.

Due to the change of operator ordering, genuine multi-point functions are, in general, discontinuous, when ever two contour coordinates become identical. In particular, two-point functions like $iF(x, y) = \langle \mathcal{T}_C \hat{A}(x) \hat{B}(y) \rangle$ become⁵

$$iF(x, y) = \begin{pmatrix} iF^{--}(x, y) & iF^{-+}(x, y) \\ iF^{+-}(x, y) & iF^{++}(x, y) \end{pmatrix} = \begin{pmatrix} \langle \mathcal{T} \hat{A}(x) \hat{B}(y) \rangle & \mp \langle \hat{B}(y) \hat{A}(x) \rangle \\ \langle \hat{A}(x) \hat{B}(y) \rangle & \langle \mathcal{T}^{-1} \hat{A}(x) \hat{B}(y) \rangle \end{pmatrix} \tag{A5}$$

where $\mathcal{T}$ and $\mathcal{T}^{-1}$ are the usual time and anti-time ordering operators. Since there are altogether only two possible orderings of the two operators, in fact given by the Wightman functions F^{-+} and F^{+-} , which are both continuous, not all four components of F are independent. Eq. (A5) implies the following relations between nonequilibrium and usual retarded and advanced functions

$$\begin{aligned}
F^R(x, y) &= F^{--}(x, y) - F^{-+}(x, y) = F^{+-}(x, y) - F^{++}(x, y) \\
&:= \Theta(x_0 - y_0) (F^{+-}(x, y) - F^{-+}(x, y)), \\
F^A(x, y) &= F^{--}(x, y) - F^{+-}(x, y) = F^{-+}(x, y) - F^{++}(x, y) \\
&:= -\Theta(y_0 - x_0) (F^{+-}(x, y) - F^{-+}(x, y)),
\end{aligned} \tag{A6}$$

⁵ Frequently used alternative notation is $F^< = F^{-+}$ and $F^> = F^{+-}$.

where $\Theta(x_0 - y_0)$ is the step function of the time difference. The rules for the co-contour functions F_{--} etc. follow from Eq. (A3).

For such two point functions complex conjugation implies

$$\begin{aligned}
(iF^{-+}(x, y))^* &= iF^{-+}(y, x) \quad \Rightarrow \quad iF^{-+}(X, p) = \text{real}, \\
(iF^{+-}(x, y))^* &= iF^{+-}(y, x) \quad \Rightarrow \quad iF^{+-}(X, p) = \text{real}, \\
(iF^{--}(x, y))^* &= iF^{++}(y, x) \quad \Rightarrow \quad (iF^{--}(X, p))^* = iF^{++}(X, p), \\
(F^R(x, y))^* &= F^A(y, x) \quad \Rightarrow \quad (F^R(X, p))^* = F^A(X, p),
\end{aligned} \tag{A7}$$

where the right parts specify the corresponding properties in the Wigner representation. Diagrammatically these rules imply the simultaneous swapping of all $+$ vertices into $-$ vertices and vice versa together with reversing the line arrow-sense of all propagator lines in the diagram.

Contrary to the common case (A5), the symmetrized contour convolution

$$E(x) = \int_{\mathcal{C}} dz [F(x, z)G(z, x) + G(x, z)F(z, x)] \tag{A8}$$

is continuous, when two contour coordinates become identical. This can be easily checked, proceeding from relations (A6) for F and G functions. Moreover, for this symmetrized convolution with two coincident points we obtain a very simple expression in the Wigner representation, if all gradient corrections to the convolution are neglected (so-called local approximation),

$$E^{\text{loc}}(X) = \int \frac{d^4 p}{(2\pi)^4} [F^{--}(X, p)G^{--}(X, p) - F(X, p)^{++}G^{++}(X, p)]. \tag{A9}$$

In particular, this form is applicable to the potential (25) and derivative (D4) energy densities.

Appendix B: DERIVATIVE COUPLING

To be specific, we consider a multicomponent system with different constituents a described by nonrelativistic fermionic and relativistic scalar bosonic field operators, summa-

rized as $\widehat{\phi} = \{\widehat{\phi}_a(x)\}$. The free Lagrangians of these fields are

$$\widehat{\mathcal{L}}_a^0 = \begin{cases} \frac{1}{2} \left(i\widehat{\phi}_a^\dagger \partial_t \widehat{\phi}_a - i\partial_t \widehat{\phi}_a^\dagger \widehat{\phi}_a - \frac{1}{m_a} \nabla \widehat{\phi}_a^\dagger \nabla \widehat{\phi}_a \right) & \text{nonrel. fermions} \\ \frac{1}{2} \left(\partial_\mu \widehat{\phi}_a \cdot \partial^\mu \widehat{\phi}_a - m_a^2 \widehat{\phi}_a^2 \right) & \text{neutral rel. bosons} \\ \partial_\mu \widehat{\phi}_a^\dagger \partial^\mu \widehat{\phi}_a - m_a^2 \widehat{\phi}_a^\dagger \widehat{\phi}_a & \text{charged rel. bosons} \end{cases} \quad (\text{B1})$$

We assume that these fields interact via linear derivative coupling, such that the interaction Lagrangian does not only depend on these fields but also on their derivatives $\widehat{\mathcal{L}}^{\text{int}} = \widehat{\mathcal{L}}^{\text{int}}\{\widehat{\phi}_a, \widehat{\phi}_a^\dagger, \partial^\mu \widehat{\phi}_a, \partial^\mu \widehat{\phi}_a^\dagger\}$. The variational principle of stationary action determines Euler–Lagrange equations of motion for the field operators

$$\partial_\mu \frac{\partial \widehat{\mathcal{L}}^0}{\partial (\partial_\mu \widehat{\phi}_a^\dagger)} - \frac{\partial \widehat{\mathcal{L}}^0}{\partial (\widehat{\phi}_a^\dagger)} = \frac{\partial \widehat{\mathcal{L}}^{\text{int}}}{\partial (\widehat{\phi}_a^\dagger)} - \partial_\mu \frac{\partial \widehat{\mathcal{L}}^{\text{int}}}{\partial (\partial_\mu \widehat{\phi}_a^\dagger)} =: \frac{\delta \widehat{\mathcal{L}}^{\text{int}}}{\delta \widehat{\phi}_a^\dagger(x)} \quad (\text{B2})$$

and the corresponding adjoint equation, where the “variational” δ -derivative is defined as

$$\frac{\delta}{\delta f(x)} \cdots := \frac{\partial}{\partial f(x)} \cdots - \partial_\mu \left(\frac{\partial}{\partial (\partial_\mu f(x))} \cdots \right). \quad (\text{B3})$$

This is the key definition, which allows to recast the local-coupling formulas to the derivative coupling case. In fact, the “variational” δ -derivative specifies the *full* derivative over $f(x)$, implying that all derivatives acting on $f(x)$ in the action should be redirected to other terms by means of partial integration before taking variational derivatives of $f(x)$.

The equations of motion can also be written in the differential form

$$G_0^{-1}(x) \widehat{\phi}_a(x) = -\widehat{J}_a(x) \equiv -\frac{\delta \widehat{\mathcal{L}}^{\text{int}}}{\delta \widehat{\phi}_a^\dagger(x)} \quad (\text{B4})$$

and similarly for the corresponding adjoint equation. The $\widehat{J}_a(x)$ operator is the local source current of field a with mass m_a , and $G_0^{-1}(x)$ is the free evolution operator⁶

$$G_0^{-1}(x) = \begin{cases} -\partial_\mu \partial^\mu - m_a^2 & \text{for rel. bosons} \\ i\partial_t + \frac{1}{2m_a} \partial_x^2 & \text{for nonrel. particles} \end{cases} \quad (\text{B5})$$

with free propagator $G_0(y, x)$ as resolvent, cf. Eq. (2).

⁶ Note that the first line in (B5) is not the nonrelativistic limit of the second one. We have already divided the second line by $2m_a$, to take into account different normalizations of relativistic and nonrelativistic wave functions.

Invariances of the Lagrangian provide a set of conservation laws, the most prominent of which are those for the energy–momentum and certain currents. In addition to the standard canonical energy–momentum tensor [28], different representations of this tensor have been considered [29, 30]. Using the Euler–Lagrange equations of motion and the definition of the source current (B4), one can show that the following form also defines a conserving energy–momentum tensor

$$\begin{aligned} \hat{\Theta}^{\mu\nu}(x) = & -\frac{1}{2} \left[\sum_a \left(\frac{1}{2}\right)_{\text{n.b.}} (\partial_x^\nu - \partial_y^\nu) \left(\frac{\partial \hat{\mathcal{L}}^0(x)}{\partial (\partial_\mu \hat{\phi}_a)} \hat{\phi}_a(y) - \hat{\phi}_a^\dagger(x) \frac{\partial \hat{\mathcal{L}}^0(y)}{\partial (\partial_\mu \hat{\phi}_a^\dagger)} \right) \right]_{x=y} \\ & + g^{\mu\nu} \left(\hat{\mathcal{E}}^{\text{int}}(x) - \hat{\mathcal{E}}^{\text{pot}}(x) \right) + \hat{\mathcal{E}}_{(\text{der.})}^{\mu\nu}(x). \end{aligned} \quad (\text{B6})$$

Here we have introduced the operators of the interaction-energy density $\hat{\mathcal{E}}^{\text{int}}$ and the potential-energy density $\hat{\mathcal{E}}^{\text{pot}}$

$$\hat{\mathcal{E}}^{\text{int}}(x) = -\hat{\mathcal{L}}^{\text{int}}(x), \quad (\text{B7})$$

$$\hat{\mathcal{E}}^{\text{pot}}(x) = -\frac{1}{2} \sum_a \left(\frac{1}{2}\right)_{\text{n.b.}} \left(\hat{J}_a^\dagger(x) \hat{\phi}_a(x) + \hat{\phi}_a^\dagger(x) \hat{J}_a(x) \right). \quad (\text{B8})$$

Furthermore, we have singled out the contribution

$$\hat{\mathcal{E}}_{(\text{der.})}^{\mu\nu}(x) = \sum_a \left(\frac{1}{2}\right)_{\text{n.b.}} \left(\frac{\partial \hat{\mathcal{L}}^{\text{int}}}{\partial (\partial_\mu \hat{\phi}_a)} \cdot \partial^\nu \hat{\phi}_a + \partial^\nu \hat{\phi}_a^\dagger \cdot \frac{\partial \hat{\mathcal{L}}^{\text{int}}}{\partial (\partial_\mu \hat{\phi}_a^\dagger)} \right) \quad (\text{B9})$$

arising in the case of derivative coupling. Here and below the case of neutral bosons results from equating $\hat{\phi}_a = \hat{\phi}_a^\dagger$ in all the formulas. Proper counting is assured by the extra $\left(\frac{1}{2}\right)_{\text{n.b.}}$ factor which takes the value 1/2 for neutral boson (real fields) and 1 for complex fields.

If the Lagrangian is invariant under some global transformation of charged fields (with the charges e_a), e.g.,

$$\hat{\phi}_a(x) \Rightarrow e^{-ie_a\Lambda} \hat{\phi}_a(x); \quad \hat{\phi}_a^\dagger(x) \Rightarrow e^{ie_a\Lambda} \hat{\phi}_a^\dagger(x), \quad (\text{B10})$$

there exists a Noether current defined as [28]

$$\hat{j}^\mu = -i \sum_a e_a \left(\frac{\partial \hat{\mathcal{L}}}{\partial (\partial_\mu \hat{\phi}_a)} \hat{\phi}_a - \hat{\phi}_a^\dagger \frac{\partial \hat{\mathcal{L}}}{\partial (\partial_\mu \hat{\phi}_a^\dagger)} \right) = \hat{j}_{(\text{conv.})}^\mu + \hat{j}_{(\text{der.})}^\mu. \quad (\text{B11})$$

which is conserved, i.e. $\partial_\mu \hat{j}^\mu = 0$. Here, we have decomposed it into two terms: the conventional one

$$\hat{j}_{(\text{conv.})}^\mu = -i \sum_a e_a \left(\frac{\partial \hat{\mathcal{L}}^0}{\partial (\partial_\mu \hat{\phi}_a)} \hat{\phi}_a - \hat{\phi}_a^\dagger \frac{\partial \hat{\mathcal{L}}^0}{\partial (\partial_\mu \hat{\phi}_a^\dagger)} \right), \quad (\text{B12})$$

which is associated with the free Lagrangian, and the derivative term

$$\hat{j}_{(\text{der.})}^\mu = -i \sum_a e_a \left(\frac{\partial \hat{\mathcal{L}}^{\text{int}}}{\partial (\partial_\mu \hat{\phi}_a)} \hat{\phi}_a - \hat{\phi}_a^\dagger \frac{\partial \hat{\mathcal{L}}^{\text{int}}}{\partial (\partial_\mu \hat{\phi}_a^\dagger)} \right) \quad (\text{B13})$$

which is non-zero only for derivative coupling.

To define the Φ -functional for the case under consideration, all the steps described in ref. [17] should be repeated. Then we arrive to the Φ -functional that depends also on the gradients of mean fields ($\partial_\mu \phi_a$ and $\partial_\mu \phi_a^*$) and Green functions ($\partial_x^\mu G(x, y)$ and $\partial_y^\mu G(x, y)$) rather than on their values only. The variational rules of this functional formally look similarly to those in ref. [17]

$$iJ_a(x) = \frac{\delta i\Phi}{\delta \phi_a^*(x)}, \quad (\text{B14})$$

$$-i\Sigma_a(x, y) = (\mp) \frac{\delta i\Phi}{\delta iG_a(y, x)} \times \begin{cases} 2 & \text{for real fields} \\ 1 & \text{for complex fields} \end{cases} \quad (\text{B15})$$

$$-\mathcal{E}^{\text{int}}(x) = \frac{\delta i\Phi}{\delta i\lambda(x)}, \quad (\text{B16})$$

but should be understood in terms of the variational δ -derivative of Eq. (B3) for one-point functions (like $\phi_a(x)$ and $\lambda(x)$) and its generalization

$$\begin{aligned} \frac{\delta i\Phi}{\delta iG(y, x)} &:= \frac{\delta_0 i\Phi}{\delta_0 iG(y, x)} - \partial_x^\mu \left(\frac{\delta_0 i\Phi}{\delta_0 (\partial_x^\mu iG(y, x))} \right) - \partial_y^\mu \left(\frac{\delta_0 i\Phi}{\delta_0 (\partial_y^\mu iG(y, x))} \right) \\ &\quad + \partial_x^\mu \partial_y^\nu \left(\frac{\delta_0 i\Phi}{\delta_0 (\partial_x^\mu \partial_y^\nu iG(y, x))} \right) \end{aligned} \quad (\text{B17})$$

to two-point functions. Here $\delta_0/\delta_0 iG(y, x)$ means the conventional variation over $G(y, x)$, which does not touch $\partial_x^\mu iG$, $\partial_y^\mu iG$ and $\partial_x^\mu \partial_y^\nu iG$ terms in $i\Phi$. Similarly to the variational δ -derivative of Eq. (B3), the δ -variation of Eq. (B17) means the *full* variation over $G(y, x)$, implying that all derivatives acting on $G(y, x)$ in the Φ functional should be redirected to other terms by means of partial integration before taking variation over $G(y, x)$. The factor $\lambda(x)$, appeared in Eq. (B16), is an auxiliary scaling parameter of the coupling constant.

In terms of the Φ functional the additional derivative contributions to mean values of the energy–momentum tensor (B9) and current (B13) take the following form

$$\mathcal{E}_{(\text{der.})}^{\mu\nu} =: \langle \hat{\mathcal{E}}_{(\text{der.})}^{\mu\nu} \rangle = \sum_a \left(\left(\frac{1}{2} \right)_{\text{n.b.}} \left[\frac{\delta\Phi}{\delta(\partial_\mu\phi_a(x))} \partial^\nu\phi_a(x) + \frac{\delta\Phi}{\delta(\partial_\mu\phi_a^*(x))} \partial^\nu\phi_a^*(x) \right] + \int_c dz \left[\frac{\delta\Phi}{\delta(\partial_\mu^x iG_a(z, x))} \cdot \partial_x^\nu iG_a(z, x) + \partial_x^\nu iG_a(x, z) \cdot \frac{\delta\Phi}{\delta(\partial_\mu^x iG_a(x, z))} \right] \right), \quad (\text{B18})$$

$$j_{(\text{der.})}^\mu =: \langle \hat{j}_{(\text{der.})}^\mu \rangle = -i \sum_a e_a \left(\left[\frac{\delta\Phi}{\delta(\partial_\mu\phi_a(x))} \phi_a(x) - \frac{\delta\Phi}{\delta(\partial_\mu\phi_a^*(x))} \phi_a^*(x) \right] + \int_c dz \left[iG_a(x, z) \cdot \frac{\delta\Phi}{\delta(\partial_\mu^x iG_a(x, z))} - \frac{\delta\Phi}{\delta(\partial_\mu^x iG_a(z, x))} \cdot iG_a(z, x) \right] \right), \quad (\text{B19})$$

while the rest terms of $\Theta^{\mu\nu}$ and j^μ retain the same form as that for local coupling, cf. ref. [17]. Here the variation is also understood in terms of Eq. (B17) to take account of the $\partial_x^\mu \partial_y^\nu iG_a(y, x)$ dependence of the Φ functional.

The next step to the kinetic description consists in gradient expansion of Kadanoff–Baym equations and all the related quantities. Expansion of the equations of motion up to the first order in gradients implies that the conserving quantities and self-energies, except for possible memory terms in the collision integral, are required only up to zero order in gradients. These zero-order quantities are determined by the local Φ functional, where all gradient corrections are neglected. Since in the local approximation $\partial_x^\mu iG$, $\partial_y^\nu iG$ and $\partial_x^\mu \partial_y^\nu iG$ transform into $-iq^\mu iG(X, q)$, $iq^\nu iG(X, q)$ and $-iq^\mu iq^\nu iG(X, q)$, respectively, no partial integrations are needed for the variations of Eqs. (B15)–(B16). This means that conventional variation rules of Eq. (3) still hold in this case. At the same time, derivative contributions to the conserving quantities, Eqs. (B18) and (B19), involve only variations over derivatives of the Green functions and, hence, should be carefully defined within the local approximation for the particular application considered.

Appendix C: NONRELATIVISTIC NUCLEAR MATTER

Currently, calculations of ground-state and low-temperature properties of nuclear matter are performed within the G - or T -matrix approximations to the self-energy [31, 32, 33]. Based on realistic nonrelativistic nucleon–nucleon potentials, they quantitatively reproduce phenomenological properties of nuclear matter. However, already for the ground state the

resulting chemical potential, i.e. the single particle separation energy, deviates from the binding energy per particle violating the Hugenholtz–van-Hove theorem. This is a manifestation of problems with the thermodynamic consistency in these approximations, which gets even worse at nonzero temperatures. This problem was discussed in refs. [31, 34, 35]. A consistent way to overcome this problem consists in using a self-consistent T -matrix approximation [31] based on the Φ -derivable approximation.

Dynamic simulations of the nuclear matter are even more demanding to the choice of approximation to the self-energy, because the requirement of charge and energy–momentum conservations should be met except for that of the thermodynamic consistency. Again all these requirements are met provided the approximation is Φ -derivable. Since dynamic simulations are much more complicated as compared to static ones, up to now they were performed in a simpler approximation to the self-energy, i.e. the direct Born approximation [5, 7], which provide qualitative description of the dynamics. These simulations were based on the Kadanoff–Baym equations without any gradient expansion. Here we would like to call attention to the fact that the use of the gradient expansion in the KB form (see subsect. IIC) would simplify these dynamic simulations and at the same time preserve the pleasant features of exact conservations and thermodynamic consistency.

In view of reasonable level of complexity feasible for current computing, we confine our consideration to the full Born approximation to the Φ functional

$$i\Phi^{\text{HFB}} = \underbrace{\frac{1}{2} \text{ (two self-energy loops)} + \frac{1}{2} \text{ (exchange self-energy)}}_{\Phi^{\text{HF}}} + \underbrace{\frac{1}{4} \text{ (two-particle exchange)} + \frac{1}{4} \text{ (two-particle exchange with crossed lines)}}_{\Phi^{\text{Born}}} \quad (\text{C1})$$

which includes the Hartree–Fock contribution Φ^{HF} (the first two diagrams in Eq. (C1)) and the true Born contribution Φ^{Born} (the last two diagrams). Here the wave line symbolizes a nonlocal nucleon–nucleon potential $V(|\mathbf{x}_1 - \mathbf{x}_2|)$, or $V(|\mathbf{q}|)$ in the momentum representation. For simplicity, below we denote the latter as $V(q)$, keeping in mind that in fact it does not depend on neither q_0 nor direction of $\mathbf{q}$.

Note that the Φ^{Born} part gives rise to the self-energy containing internal vertices. This implies that the corresponding collision term involves non-local effects (see discussion in ref. [8]). However, only “spatial non-locality” appears in the collision term, while the memory in time is absent since $V(|\mathbf{x}_1 - \mathbf{x}_2|)\delta(t_1 - t_2)$ is time local. According to the general consideration of ref. [10], exact conservations in the gradient approximation take place if all the non-local terms are consistently taken into account up to first-order gradients. Below we show that in the particular case of the Φ^{HFB} functional the exact conservations hold true even if we neglect the “spatial non-locality” generated by Φ^{HFB} . These exact conservations imply the global rather than local conservation of the energy-momentum, which is in fact natural for the case of instant interaction of finite range considered here.

Neglecting the gradient terms induced by the finite range of V , we consider the Φ^{Born} functional in the local approximation, where all Green functions in the Wigner representation are taken at the same centroid coordinate X . Alongside in some variational expressions we use an X -dependent local Φ functional, $\Phi(X)$, where the last spatial integration is omitted, i.e.

$$\Phi = \int dX \Phi(X) \quad (\text{C2})$$

The Φ^{HFB} of Eq. (C1) gives rise to the following local collision term

$$C^{\text{HFB-loc}} = \int \frac{d^4 p_2}{(2\pi)^4} \frac{d^4 p_3}{(2\pi)^4} \frac{d^4 p_4}{(2\pi)^4} R^{\text{HFB}} \times \left(\tilde{F}_1 \tilde{F}_2 F_3 F_4 - F_1 F_2 \tilde{F}_3 \tilde{F}_4 \right) \delta^4(p_1 + p_2 - p_3 - p_4), \quad (\text{C3})$$

$$R^{\text{HFB}} = \frac{(2\pi)^4}{2} [V(p_1 - p_3) + V(p_1 - p_4)]^2, \quad (\text{C4})$$

where $F_1 = F(X, p_1)$, etc., cf. Eq. (17).

1. Charge Conservation

In Φ^{HFB} the G^{--} and G^{++} Green functions are encountered only in different $+-$ Φ -diagrams, and hence we can vary G^{--} and G^{++} independently. Therefore, Φ^{HFB} is invariant under the following transformation

$$G^{--}(X, p) \Rightarrow G^{--}(X, p + \xi(X)), \quad G^{++}(X, p) \Rightarrow G^{++}(X, p - \xi(X)) \quad (\text{C5})$$

with F , $\tilde{F}$ and V kept unchanged. Here $\xi(X)$ is an arbitrary function. If $|\xi(X)| \ll 1$, transformation (C5) reads

$$\delta G^{--} = \xi_\mu(X) \frac{\partial G^{--}}{\partial p_\mu}, \quad \delta G^{++} = -\xi_\mu(X) \frac{\partial G^{++}}{\partial p_\mu}. \quad (\text{C6})$$

Performing variation of Φ^{Born} under the transformation (C6) within the canonical variation rules (3), we arrive at

$$\begin{aligned} i\delta\Phi^{\text{loc}} &= \int dX \xi_\mu(X) \text{Tr} \int \frac{d^4p}{(2\pi)^4} \left(i\Sigma_{--} \frac{\partial iG^{--}}{\partial p_\mu} - i\Sigma_{++} \frac{\partial iG^{++}}{\partial p_\mu} \right) \\ &= 2i \int dX \xi_\mu(X) \text{Tr} \int \frac{d^4p}{(2\pi)^4} \left(\Gamma^{\text{in}} \frac{\partial \text{Re } G^R}{\partial p_\mu} + \text{Re } \Sigma^R \frac{\partial F}{\partial p_\mu} \right) = 0. \end{aligned} \quad (\text{C7})$$

Here we have used that the integral

$$\text{Tr} \int \frac{d^4p}{(2\pi)^4} \left(\Gamma \frac{\partial \text{Re } G^R}{\partial p_\mu} + \text{Re } \Sigma^R \frac{\partial A}{\partial p_\mu} \right) = -\frac{1}{2} \text{Im} \text{Tr} \int \frac{d^4p}{(2\pi)^4} \Sigma^R \frac{\partial G^R}{\partial p_\mu} = 0. \quad (\text{C8})$$

equals zero due to analyticity of G^R and Σ^R . Thus, we obtain the relation

$$\text{Tr} \int \frac{d^4p}{(2\pi)^4} \left(\Gamma^{\text{in}} \frac{\partial \text{Re } G^R}{\partial p_\mu} + \text{Re } \Sigma^R \frac{\partial F}{\partial p_\mu} \right) = 0, \quad (\text{C9})$$

which guarantees the Noether current conservation, cf. Eq. (21) with $j_{(\text{der})}^\mu = 0$.

2. Energy–Momentum Conservation

In order to construct the conservation laws related to space–time homogeneity we perform the following transformation

$$G^{--}(X, p) \Rightarrow G^{--}(X + \xi(X), p), \quad G^{++}(X, p) \Rightarrow G^{++}(X - \xi(X), p) \quad (\text{C10})$$

with F , $\tilde{F}$ and V kept unchanged. This transformation only acts on $\Phi^{\text{HFB}-}$ and $\Phi^{\text{HFB}+}$, i.e. those involving only “−” or “+” vertices, respectively,

$$\delta\Phi^{\text{HFB}} = \int dX \xi_\mu(X) \partial^\mu (\Phi^{\text{HFB}}(X^-) - \Phi^{\text{HFB}}(X^+)) \quad (\text{C11})$$

where $\Phi^{\text{HFB}}(X^i)$ are understood in the sense of (C2). Note that

$$\begin{aligned} i(\Phi^{\text{HFB}}(X^-) - \Phi^{\text{HFB}}(X^+)) &= i\text{Tr} \int \frac{d^4p}{(2\pi)^4} \Sigma^{\text{HF}} F \\ &+ \frac{1}{2} i\text{Tr} \int \frac{d^4p}{(2\pi)^4} (\Gamma^{\text{in}} \text{Re } G^R + (\text{Re } \Sigma^R - \Sigma^{\text{HF}}) F) \\ &- \underbrace{\text{Tr} \int \frac{d^4p}{(2\pi)^4} (\Gamma \text{Re } G^R + (\text{Re } \Sigma^R - \Sigma^{\text{HF}}) A)}_{=0}, \end{aligned} \quad (\text{C12})$$

where the last integral is again zero due to analyticity, similarly to (C8). Here the first term on the r.h.s. results from the first two (Hartree–Fock) diagrams in Eq. (C1), while the last two integrals follow from the last two (Born) diagrams. Alternatively, we can perform variation of Φ^{Born} applying the canonical variation rules (3)

$$\begin{aligned} i\delta\Phi^{\text{HFB}} &= \int dX \xi_\mu(X) \text{Tr} \int \frac{d^4p}{(2\pi)^4} \left(i\Sigma_{--} \frac{\partial iG^{--}}{\partial X_\mu} - i\Sigma_{++} \frac{\partial iG^{++}}{\partial X_\mu} \right) \\ &= 2i \int dX \xi_\mu(X) \text{Tr} \int \frac{d^4p}{(2\pi)^4} \left(\Gamma^{\text{in}} \frac{\partial \text{Re } G^R}{\partial X_\mu} + \text{Re } \Sigma^R \frac{\partial F}{\partial X_\mu} \right) \\ &\quad - i \underbrace{\int dX \xi_\mu(X) \text{Tr} \int \frac{d^4p}{(2\pi)^4} \left(\Gamma \frac{\partial \text{Re } G^R}{\partial X_\mu} + \text{Re } \Sigma^R \frac{\partial A}{\partial X_\mu} \right)}_{=0}, \end{aligned} \quad (\text{C13})$$

where the last integral is again zero due to analyticity. Therefore, we arrive at the important identity

$$\begin{aligned} &\text{Tr} \int \frac{d^4p}{(2\pi)^4} \left(\Gamma^{\text{in}} \frac{\partial \text{Re } G^R}{\partial X_\mu} + \text{Re } \Sigma^R \frac{\partial F}{\partial X_\mu} \right) \\ &= \partial^\mu \text{Tr} \int \frac{d^4p}{(2\pi)^4} \left[\frac{1}{2} \Sigma^{\text{HF}} F + \frac{1}{4} (\Gamma^{\text{in}} \text{Re } G^R + (\text{Re } \Sigma^R - \Sigma^{\text{HF}}) F) \right]. \end{aligned} \quad (\text{C14})$$

Next we investigate the following transformation

$$G^{--}(X, p) \Rightarrow G^{--}(X, \Lambda_{\mu\nu}(X) p^\nu), \quad G^{++}(X, p) \Rightarrow G^{++}(X, \Lambda_{\mu\nu}^{-1}(X) p^\nu) \quad (\text{C15})$$

$$V^-(q) \Rightarrow V^-(\Lambda_{\mu\nu}(X) q^\nu), \quad V^+(q) \Rightarrow V^+(\Lambda_{\mu\nu}^{-1}(X) q^\nu) \quad (\text{C16})$$

for the entire Φ^{HFB} , while $F, \tilde{F}$ are kept unchanged. For the Hartree–Fock part Φ^{HF} one finds

$$\tilde{p}_\mu = \Lambda_{\mu\nu} p^\nu, \quad \tilde{p}'_\mu = \Lambda_{\mu\nu} p'^\nu \quad \Rightarrow \quad d^4p \, d^4p' = (\det \Lambda)^{-2} d^4\tilde{p} \, d^4\tilde{p}', \quad (\text{C17})$$

$$\delta\Phi^{\text{HF}-} = \int dX [(\det \Lambda)^{-2} - 1] \Phi^{\text{HF}}(X^-), \quad (\text{C18})$$

$$\delta\Phi^{\text{HF}+} = \int dX [(\det \Lambda)^2 - 1] \Phi^{\text{HF}}(X^+), \quad (\text{C19})$$

where again $\Phi^{\text{HF}-}$ and $\Phi^{\text{HF}+}$ are the Φ^{HF} diagrams involving only "–" or "+" vertices, respectively. In general, an arbitrary diagram $\Phi^{n_G, n_\lambda(-)}$, consisting of n_G Green functions, n_λ of "–" interactions and no "+" interactions, transforms as

$$\delta\Phi^{n_G, n_\lambda(-)} = \int dX [(\det \Lambda)^{-(n_G - n_\lambda + 1)} - 1] \Phi^{n_G, n_\lambda(-)}(X), \quad (\text{C20})$$

since the change of each momentum integration gives $(\det \Lambda)^{-1}$ (cf. Eq. (C17)), while the transformation of the $\delta(p)$ -function at each vertex (apart from one vertex due to global momentum conservation of the diagram) produces $\det \Lambda$

$$\delta^4(p + p' - q) = \det \Lambda \delta^4(\tilde{p} + \tilde{p}' - \tilde{q}). \quad (\text{C21})$$

Similarly,

$$\delta \Phi^{n_G, n_\lambda(+)} = \int dX [(\det \Lambda)^{n_G - n_\lambda + 1} - 1] \Phi^{n_G, n_\lambda(+)}(X). \quad (\text{C22})$$

According to these rules the "−" and "+" diagrams of the second order in the interaction, $\Phi^{\text{Born}-}$ and $\Phi^{\text{Born}+}$, are transformed as follows

$$\delta \Phi^{\text{Born}-} = \int dX [(\det \Lambda)^{-3} - 1] \Phi^{\text{Born}-}(X), \quad (\text{C23})$$

$$\delta \Phi^{\text{Born}+} = \int dX [(\det \Lambda)^3 - 1] \Phi^{\text{Born}+}(X). \quad (\text{C24})$$

If the Λ -transformation is infinitesimal, $\Lambda_{\mu\nu}(X) = 1 + \omega_{\mu\nu}(X)$ with $|\omega_{\mu\nu}| \ll 1$ and $\det \Lambda = 1 + \text{Tr} \omega$, $\det(\Lambda^{-1}) = 1 - \text{Tr} \omega$, we obtain

$$\begin{aligned} \delta (\Phi^{\text{HFB}-} + \Phi^{\text{HFB}+}) = & - \int dX 2 \text{Tr} \omega (\Phi^{\text{HF}-}(X) - \Phi^{\text{HF}+}(X)) \\ & - \int dX 3 \text{Tr} \omega (\Phi^{\text{Born}-}(X) - \Phi^{\text{Born}+}(X)), \end{aligned} \quad (\text{C25})$$

cf. Eq. (C12). The $\Phi^{\text{Born}-+}$ and $\Phi^{\text{Born}+-}$ components, i.e. those containing both "−" and "+" vertices, are modified by only the V transformation. Moreover, this transformation leaves them invariant

$$\begin{aligned} i\delta \Phi^{\text{Born}-+} &= \int dX \omega_{\mu\nu} \text{Tr} \int \frac{d^4 p}{(2\pi)^4} p^\nu \left(\frac{\delta i \Phi^{\text{Born}-+}(X)}{\delta i V^-} \frac{\partial i V^-}{\partial p_\mu} - \frac{\delta i \Phi^{\text{Born}-+}(X)}{\delta i V^+} \frac{\partial i V^+}{\partial p_\mu} \right) \\ &= \int dX \omega_{\mu\nu} \text{Tr} \int \frac{d^4 p}{(2\pi)^4} p^\nu \left(\frac{\delta i \Phi^{\text{Born}-+}(X)}{\delta V^-} V^- - \frac{\delta i \Phi^{\text{Born}-+}(X)}{\delta V^+} V^+ \right) \frac{1}{V} \frac{\partial V}{\partial p_\mu} = 0, \end{aligned} \quad (\text{C26})$$

since $\Phi^{\text{Born}-+}$ is symmetric with respect to V^- and V^+ . Thus,

$$\begin{aligned} \delta i \Phi^{\text{HFB}} &= - \int dX (\text{Tr} \omega) i \text{Tr} \int \frac{d^4 p}{(2\pi)^4} [2 \Sigma^{\text{HF}} F \\ &\quad - \frac{3}{2} (\Gamma^{\text{in}} \text{Re} G^R + (\text{Re} \Sigma^R - \Sigma^{\text{HF}}) F)]. \end{aligned} \quad (\text{C27})$$

Alternatively, we can perform variation of Φ^{HFB} applying the canonical variation rules

(3)

$$i\delta \Phi^{\text{HFB}} = \int dX \omega_{\mu\nu} \left[\text{Tr} \int \frac{d^4 p}{(2\pi)^4} p^\nu \left(i \Sigma_{--} \frac{\partial i G^{--}}{\partial p_\mu} - i \Sigma_{++} \frac{\partial i G^{++}}{\partial p_\mu} \right) + 2i Q^{\mu\nu}(X) \right]$$

$$\begin{aligned}
&= 2i \int dX \omega_{\mu\nu} \left[\text{Tr} \int \frac{d^4 p}{(2\pi)^4} p^\nu \left(\Gamma^{\text{in}} \frac{\partial \text{Re } G^R}{\partial p_\mu} + \text{Re } \Sigma^R \frac{\partial F}{\partial p_\mu} \right) + Q^{\mu\nu}(X) \right] \\
&\quad - i \underbrace{\int dX \omega_{\mu\nu} \text{Tr} \int \frac{d^4 p}{(2\pi)^4} p^\nu \left(\Gamma \frac{\partial \text{Re } G^R}{\partial p_\mu} + \text{Re } \Sigma^R \frac{\partial A}{\partial p_\mu} \right)}_{=0}, \tag{C28}
\end{aligned}$$

where the last integral again equals zero due to analyticity. Here we have introduced the quantity

$$2iQ^{\mu\nu}(X) = \text{Tr} \int \frac{d^4 p}{(2\pi)^4} p^\nu \left(\frac{\delta i\Phi^{\text{HFB}^-}(X)}{\delta iV^-} \frac{\partial V^-}{\partial p_\mu} - \frac{\delta i\Phi^{\text{HFB}^+}(X)}{\delta iV^+} \frac{\partial V^+}{\partial p_\mu} \right) \tag{C29}$$

arising from variation over V . All we have to know about this quantity is that it is $Q^{\mu\nu} = 0$ when $\mu = 0$ and/or $\nu = 0$. This property results from p_0 -independence of $V(|\mathbf{p}|)$. In particular this property yields

$$\int d^3 X \partial_\mu Q^{\mu\nu}(X) = 0, \tag{C30}$$

where the X -integration runs only over space.

Hence, comparing Eq. (C28) to Eq. (C27), we arrive at another important identity

$$\begin{aligned}
&\text{Tr} \int \frac{d^4 p}{(2\pi)^4} p^\nu \left(\Gamma^{\text{in}} \frac{\partial \text{Re } G^R}{\partial p_\mu} + \text{Re } \Sigma^R \frac{\partial F}{\partial p_\mu} \right) + Q^{\mu\nu}(X) \\
&= -g^{\mu\nu} \text{Tr} \int \frac{d^4 p}{(2\pi)^4} \Sigma^{\text{HF}} F - g^{\mu\nu} \frac{3}{4} \text{Tr} \int \frac{d^4 p}{(2\pi)^4} (\Gamma^{\text{in}} \text{Re } G^R + (\text{Re } \Sigma^R - \Sigma^{\text{HF}}) F). \tag{C31}
\end{aligned}$$

We turn now to the r.h.s. of the consistency relation for energy-momentum conservation (22)

$$K^\nu = \text{Tr} \int d^3 X \frac{p^\nu d^4 p}{(2\pi)^4} [\{\text{Re } \Sigma^R, F\} - \{\text{Re } G^R, \Gamma^{\text{in}}\}] \tag{C32}$$

integrated over space, which is suitable for the global conservation. In this expression the local collision term (C3) drops out according to Eq. (16). It can be transformed by means of the identity

$$\int \frac{d^4 p}{(2\pi)^4} p^\nu \{\varphi, f\} = \int \frac{d^4 p}{(2\pi)^4} \left[\partial^\mu \left(p^\nu f \frac{\partial \varphi}{\partial p^\mu} \right) + f \partial^\nu \varphi \right], \tag{C33}$$

where φ and f are arbitrary functions, with the result

$$\begin{aligned}
K^\nu &= -\text{Tr} \int d^3 X \partial^\mu \int \frac{d^4 p}{(2\pi)^4} p^\nu \left(\text{Re } \Sigma^R \frac{\partial F}{\partial p^\mu} + \Gamma^{\text{in}} \frac{\partial \text{Re } G^R}{\partial p^\mu} \right) \\
&\quad - \text{Tr} \int d^3 X \frac{d^4 p}{(2\pi)^4} (\text{Re } \Sigma^R \partial^\nu F + \Gamma^{\text{in}} \partial^\nu \text{Re } G^R). \tag{C34}
\end{aligned}$$

Now, applying identities (C14), (C30) and (C31) to the r.h.s. of Eq. (C34), we obtain

$$K^\nu = \int d^3X \partial_\mu g^{\mu\nu} \text{Tr} \int \frac{d^4p}{(2\pi)^4} \frac{1}{2} (\Gamma^{\text{in}} \text{Re} G^R + \text{Re} \Sigma^R F) \quad (\text{C35})$$

which is precisely needed for the global conservation of the Noether energy–momentum

$$\frac{\partial}{\partial T} \int d^3X \Theta^{0\nu}(X) = 0, \quad (\text{C36})$$

since for the case under consideration $\mathcal{E}^{\text{int}} = \frac{1}{2}\mathcal{E}^{\text{pot}}$, cf. Eq. (26).

Appendix D: NUCLEON–PION SYSTEM

For the discussion of the physical aspects of the nucleon–pion problem we refer to ref. [9]. Here we would like to clarify some technical details. We choose the nonrelativistic form of pion–nucleon interaction [36]

$$\widehat{\mathcal{L}}^{\text{int}} = g \widehat{\psi}^\dagger [(\boldsymbol{\sigma} \cdot \boldsymbol{\nabla})(\boldsymbol{\tau} \cdot \widehat{\boldsymbol{\phi}})] \widehat{\psi}, \quad (\text{D1})$$

where $\widehat{\psi}$ and $\widehat{\phi}$ are non-relativistic nucleon and Klein-Gordon pion field operators, respectively. We accept a simple approximation defined by the following Φ functional

$$\begin{aligned} i\Phi^{\pi N} &= \frac{1}{2} \text{Tr} \int dX \frac{d^4p_1}{(2\pi)^4} \frac{d^4p_2}{(2\pi)^4} \frac{d^4q}{(2\pi)^4} \delta^4(p_1 - p_2 + q) \\ &\times iG^{ij}(X, p_1) (-ig)(-i\mathbf{q} \cdot \boldsymbol{\sigma}) \boldsymbol{\tau} i\Delta_{ij}(X, q) (-ig)(i\mathbf{q} \cdot \boldsymbol{\sigma}) \boldsymbol{\tau} iG^{ji}(X, p_2), \end{aligned} \quad (\text{D2})$$

where G and Δ are the nucleon and pion Green functions, respectively, and Tr runs over spin and isospin indices. Here we have assumed an isotopically symmetric system, where the pion Green functions of all isotopic charges coincide. Though this approximation is evidently oversimplified to produce quantitative results, cf. ref. [9], it is able to reproduce qualitative features of the dynamics. Moreover, this approximation is at the edge of the present computing abilities. The formal basics of the Φ -functional formalism are given in Append. B.

The charge current, defined by Eqs. (20) and (B11), relates to the baryon number conservation and hence is trivial from the point of view of the pion–nucleon interaction. Indeed, to prove the baryon number conservation we should perform transformation (C5)

with F_N , $\tilde{F}_N$ and Δ^{ij} kept unchanged. The pion Green functions Δ_{ij} are not subjected to this transformation, since pions are neutral from the point of view of baryonic charge. All the subsequent considerations are completely identical to that of the Fock diagram (the second term in Eq. (C1)) and lead to the same final result (C9), i.e. to the exact Noether current conservation.

The energy–momentum conservation is more instructive in this respect. Before proceeding to the conservation laws themselves we should define the derivative contribution to the energy–momentum tensor (B18). In our case of vanishing mean fields the pion Green function enters the Φ -functional only doubly differentiated. Therefore the expression (B18) takes the form

$$\mathcal{E}_{(\text{der.})}^{\mu\nu} = \frac{1}{2} \int_{\mathcal{C}} dz \left(\frac{\delta_0 \Phi}{\delta_0(\partial_\lambda^z \partial_\mu^x \text{i}\Delta(z, x))} \cdot \partial_\lambda^z \partial_x^\nu \text{i}\Delta(z, x) + \partial_\nu^x \partial_z^\lambda \text{i}\Delta(x, z) \cdot \frac{\delta_0 \Phi}{\delta_0(\partial_\mu^x \partial_\lambda^z \text{i}\Delta(x, z))} \right), \quad (\text{D3})$$

where δ_0 is already the conventional variation. In the Wigner representation with due regard for Eq. (A9) it transforms into

$$\begin{aligned} \mathcal{E}_{(\text{der.})}^{\mu\nu}(X) = & -\text{Tr} \int \frac{d^4 p_1}{(2\pi)^4} \frac{d^4 p_2}{(2\pi)^4} \frac{d^4 q}{(2\pi)^4} \delta^4(p_1 - p_2 + q) g^2 \\ & \times \left[\text{i}G^{--}(X, p_1) (-q^\nu \sigma^\mu) \boldsymbol{\tau} \text{i}\Delta^{--}(X, q) (\text{i}\mathbf{q} \cdot \boldsymbol{\sigma}) \boldsymbol{\tau} \text{i}G^{--}(X, p_2) \right. \\ & \left. - \text{i}G^{++}(X, p_1) (-q^\nu \sigma^\mu) \boldsymbol{\tau} \text{i}\Delta^{++}(X, q) (\text{i}\mathbf{q} \cdot \boldsymbol{\sigma}) \boldsymbol{\tau} \text{i}G^{++}(X, p_2) \right]. \end{aligned} \quad (\text{D4})$$

Contrary to usual convention, here we use Greek indices μ and ν for the components of 3-vectors in order to distinguish them from the “+−” summation indices. The potential energy density is still determined by the standard expression (25) but in terms of the sum over nucleons and pions. Within the approximation of Eq. (D2), $\mathcal{E}^{\text{pot}}$ can be alternatively expressed as

$$\mathcal{E}^{\text{pot}}(X) = \mathcal{E}_N^{\text{pot}} + \mathcal{E}_\pi^{\text{pot}} = \frac{3}{2} [\Phi^{\pi N-}(X) - \Phi^{\pi N+}(X)], \quad (\text{D5})$$

where $\Phi^{\pi N-}$ ($\Phi^{\pi N+}$) refers to the $\Phi^{\pi N}$ functional with removed integration over dX and all vertices being of “−” (“+”) type. In view of relation (26),

$$\mathcal{E}^{\text{int}}(X) - \mathcal{E}^{\text{pot}}(X) = -\frac{1}{2} [\Phi^{\pi N-}(X) - \Phi^{\pi N+}(X)]. \quad (\text{D6})$$

1. Energy–Momentum Conservation

We briefly repeat the steps proving the exact energy–momentum conservation for the nonrelativistic nuclear matter (Append. C2) with the emphasis on the specifics of the derivative coupling.

First the transformation of Eq. (C10) for the nucleon Green functions together with the corresponding transformation of the pion Green functions

$$\Delta^{--}(X, p) \Rightarrow \Delta^{--}(X + \xi(X), p), \quad \Delta^{++}(X, p) \Rightarrow \Delta^{++}(X - \xi(X), p), \quad (\text{D7})$$

with F_π and $\tilde{F}_\pi$ being kept unchanged has to be performed. This transformation is unaffected by the derivative coupling, and in the similar way as before we arrive at the identity

$$\begin{aligned} & \text{Tr} \int \frac{d^4 p}{(2\pi)^4} \left[\left(\Gamma_N^{\text{in}} \frac{\partial \text{Re } G^R}{\partial X_\mu} + \text{Re } \Sigma^R \frac{\partial F_N}{\partial X_\mu} \right) + \frac{1}{2} \left(\Gamma_\pi^{\text{in}} \frac{\partial \text{Re } \Delta^R}{\partial X_\mu} + \text{Re } \Pi^R \frac{\partial F_\pi}{\partial X_\mu} \right) \right] \\ &= -\partial^\mu (\mathcal{E}^{\text{int}} - \mathcal{E}^{\text{pot}}). \end{aligned} \quad (\text{D8})$$

Here Σ and Π refer to nucleon and pion self-energies, respectively, and subscripts N or π correspondingly attribute a quantity to either nucleon or pion subsystems. The r.h.s. of this identity is written with due regards to Eq. (D6).

Let us now perform the transformation (C15) for both nucleon and pion Green functions, as well as q factors encountered in vertices of $\Phi^{\pi N}$. Then the variation of $\Phi^{\pi N}$ gives

$$\begin{aligned} \delta \Phi^{\pi N} &= \int dX \omega_{\mu\nu} \left[-2g^{\mu\nu} (\Phi^{\pi N-}(X) - \Phi^{\pi N+}(X)) + 2\mathcal{E}_{(\text{der.})}^{\mu\nu} \right] \\ &= \int dX \omega_{\mu\nu} \left[4g^{\mu\nu} (\mathcal{E}^{\text{int}} - \mathcal{E}^{\text{pot}}) + 2\mathcal{E}_{(\text{der.})}^{\mu\nu} \right], \end{aligned} \quad (\text{D9})$$

where the $\mathcal{E}_{(\text{der.})}^{\mu\nu}$ results from the variation of q factors in vertices of $\Phi^{\pi N}$. Alternatively, performing variation of $\Phi^{\pi N}$ accordingly to the canonical variation rules (3) and equating the result to expression (D9), we arrive at another identity

$$\begin{aligned} & \text{Tr} \int \frac{d^4 p}{(2\pi)^4} \left[\left(\Gamma_N^{\text{in}} \frac{\partial \text{Re } G^R}{\partial p_\mu} + \text{Re } \Sigma^R \frac{\partial F_N}{\partial p_\mu} \right) + \frac{1}{2} \left(\Gamma_\pi^{\text{in}} \frac{\partial \text{Re } \Delta^R}{\partial p_\mu} + \text{Re } \Pi^R \frac{\partial F_\pi}{\partial p_\mu} \right) \right] \\ &= 2g^{\mu\nu} (\mathcal{E}^{\text{int}} - \mathcal{E}^{\text{pot}}) + \mathcal{E}_{(\text{der.})}^{\mu\nu}, \end{aligned} \quad (\text{D10})$$

The r.h.s. of the consistency relation for energy–momentum conservation (22) now reads

$$K^\nu = \text{Tr} \int \frac{p^\nu d^4 p}{(2\pi)^4} \left[(\{\text{Re } \Sigma^R, F_N\} - \{\text{Re } G^R, \Gamma_N^{\text{in}}\}) \right]$$

$$+ \frac{1}{2} (\{\text{Re } \Pi^R, F_\pi\} - \{\text{Re } \Delta^R, \Gamma_\pi^{\text{in}}\}) , \quad (\text{D11})$$

By means of identity (C33) it is transformed to the form

$$\begin{aligned} K^\nu = & -\partial^\mu \text{Tr} \int \frac{d^4 p}{(2\pi)^4} p^\nu \left[\left(\text{Re } \Sigma^R \frac{\partial F_N}{\partial p^\mu} + \Gamma_N^{\text{in}} \frac{\partial \text{Re } G^R}{\partial p^\mu} \right) \right. \\ & \left. + \frac{1}{2} \left(\text{Re } \Pi^R \frac{\partial F_\pi}{\partial p^\mu} + \Gamma_\pi^{\text{in}} \frac{\partial \text{Re } \Delta^R}{\partial p^\mu} \right) \right] \\ & - \text{Tr} \int \frac{d^4 p}{(2\pi)^4} \left[(\text{Re } \Sigma^R \partial^\nu F_N + \Gamma_N^{\text{in}} \partial^\nu \text{Re } G^R) \right. \\ & \left. + \frac{1}{2} (\text{Re } \Pi^R \partial^\nu F_\pi + \Gamma_\pi^{\text{in}} \partial^\nu \text{Re } \Delta^R) \right] . \end{aligned} \quad (\text{D12})$$

Now, applying identities (D8) and (D10) to the r.h.s. of Eq. (D12), we obtain

$$K^\nu = \partial^\nu (\mathcal{E}^{\text{pot}} - \mathcal{E}^{\text{int}}) - \partial_\mu \mathcal{E}_{(\text{der.})}^{\mu\nu} \quad (\text{D13})$$

which is precisely needed for the local conservation of the Noether energy–momentum.

-
- [1] S.T. Belyaev and G.I. Budker, Dokl. AN SSSR **107**, 807 (1965).
 - [2] J. Schwinger, J. Math. Phys. **2**, 407 (1961).
 - [3] L.P. Kadanoff and G. Baym, *Quantum Statistical Mechanics*, Benjamin, 1962.
 - [4] L.P. Keldysh, ZhETF **47**, 1515 (1964) [Sov. Phys. JETP **20**, 1018 (1965)].
 - [5] P. Danielewicz, Ann. Phys. (NY) **152**, 305 (1984).
 - [6] J. Knoll and D.N. Voskresensky, Phys. Lett. **B 351**, 43 (1995); Ann. Phys. (NY) **249**, 532 (1996).
 - [7] P. Bozek, Phys. Rev. C **56**, 1452 (1997).
 - [8] Yu.B. Ivanov, J. Knoll and D.N. Voskresensky, Nucl. Phys. A **672**, 313 (2000).
 - [9] Yu.B. Ivanov, J. Knoll, H. van Hees and D. N. Voskresensky, Yad. Fiz. **64**, 711 (2001).
 - [10] J. Knoll, Yu.B. Ivanov, and D.N. Voskresensky, Ann. Phys. (NY) **293**, 126 (2001).
 - [11] W. Cassing and S. Juchem, Nucl. Phys. A **665**, 377 (2000); Nucl. Phys. A **672**, 417 (2000).
 - [12] S. Leupold, Nucl. Phys. A **672**, 475 (2000).
 - [13] M. Effenberger, U. Mosel, Phys. Rev. C **60**, 51901 (1999).
 - [14] P. Danielewicz, Ann. Phys. (NY) **152**, 239 (1984).
 - [15] W. Botermans and R. Malfliet, Phys. Rep. **198**, 115 (1990).
 - [16] G. Baym, Phys. Rev. **127**, 1391 (1962).
 - [17] Yu.B. Ivanov, J. Knoll, and D.N. Voskresensky, Nucl. Phys. A **657**, 413 (1999).
 - [18] R.E. Norton and J.M. Cornwall, Ann. Phys. (NY) **92**, 106 (1975).

- [19] G.M. Carneiro and C.J. Pethick, Phys. Rev. B **11**, 1106 (1975).
- [20] G. Aarts and J. Berges, Phys. Rev. D **64**, 105010 (2001).
- [21] J. Berges, S. Borsanyi, and J. Serreau, e-Print Archive: hep-ph/0212404.
- [22] J. Berges and J. Serreau, e-Print Archive: hep-ph/0208070.
- [23] H. van Hees and J. Knoll, Phys.Rev. D **65**, 025010 (2002).
- [24] H. van Hees and J. Knoll, Phys.Rev. D **65**, 105005 (2002).
- [25] G. Baym and G. Grinstein, Phys. Rev. D **15**, 2897 (1977).
- [26] H. van Hees and J. Knoll, Phys.Rev. D **66**, 025028 (2002).
- [27] A. Arrizabalaga and J. Smit, Phys.Rev. D **66**, 065014 (2002).
- [28] C. Itzykson and J.-B. Zuber, *Quantum Field Theory*, McGraw-Hill, 1980.
- [29] J. Belinfante, Physica **6**, 887 (1939).
- [30] C.G. Callan, S. Coleman and R. Jackiw, Ann. Phys. (NY) **59**, 42 (1970).
- [31] P. Bożek and P. Czerski, Eur. Phys. J. A **11**, 271 (2001); P. Bożek, Phys. Rev. C **59**, 2619 (1999); Nucl. Phys. A **657**, 187 (1999).
- [32] W.H. Dickhoff, Phys. Rev. C **58**, 2807 (1998); W.H. Dickhoff *et al.*, Phys. Rev. C **60**, 4319 (1999).
- [33] Y. Dewulf, D. Van Neck, and M. Waroquier, , Phys. Lett. B **510**, 89 (2001).
- [34] M. Baldo *et al.*, Phys. Rev. C **41**, 1748 (1990).
- [35] F. de Jong and R. Malfliet, Phys. Rev. C **44**, 998 (1991).
- [36] T. Ericson and W. Weise, *Pions and Nuclei*, Clarendon Press, Oxford 1988; A.B. Migdal, E.E. Saperstein, M.A. Troitsky, and D.N. Voskresensky, Phys. Rep. **192**, 179 (1990).